

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/6597683>

Functional alterations in macrophages after hypoxia selection

Article in *Experimental Biology and Medicine* · February 2007

Source: PubMed

CITATIONS

22

READS

141

2 authors, including:



Selma Giorgio

University of Campinas

96 PUBLICATIONS 1,337 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Leishmaniasis investigation by MALDI Imaging Mass Spectrometry [View project](#)



Ecotoxicology and Nanotoxicology [View project](#)

Functional Alterations in Macrophages After Hypoxia Selection

Adriana Degrossoli and Selma Giorgio
Exp Biol Med (Maywood) 2007 232: 88

The online version of this article can be found at:
<http://ebm.sagepub.com/content/232/1/88>

Published by:



<http://www.sagepublications.com>

On behalf of:



Society for Experimental Biology and Medicine

Additional services and information for *Experimental Biology and Medicine* can be found at:

Email Alerts: <http://ebm.sagepub.com/cgi/alerts>

Subscriptions: <http://ebm.sagepub.com/subscriptions>

Reprints: <http://www.sagepub.com/journalsReprints.nav>

Permissions: <http://www.sagepub.com/journalsPermissions.nav>

>> [Version of Record](#) - Jan 1, 2007

[What is This?](#)

Functional Alterations in Macrophages After Hypoxia Selection

ADRIANA DEGROSSOLI AND SELMA GIORGIO¹

Department of Parasitology, Biology Institute, Universidade Estadual de Campinas, Campinas 13083-970, São Paulo, Brazil

Regions of low oxygen tension are common features of inflamed and infected tissues and provide physiologic selective pressure for the expansion of cells with enhanced hypoxia tolerance. The aim of this study was to investigate whether macrophages resistant to death induced by hypoxia were accompanied by functional alterations. A mouse macrophage cell line (J774 cells) was used to obtain subpopulations of death-resistant macrophages induced by long-term exposure to severe hypoxia (<1% O₂). The results indicated that exposing J774 macrophages to periods of severe hypoxia results in the selection of cells with phenotypes associated with the modulation of heat-shock protein 70 kDa (HSP70) expression, tumor necrosis factor- α (TNF- α), and nitric oxide (NO) production and reduced susceptibility to parasite *Leishmania* infection. Thus, we suggest that hypoxia-selected macrophages may influence the outcome of inflammation and infection. *Exp Biol Med* 232:88–95, 2007

Key words: hypoxia; selection; macrophages; inflammatory responses; *Leishmania*

Regions of low oxygen tension (hypoxia) are common features of tumors, wounds, atherosclerotic lesions, and inflamed/infected tissues (1–4). Since cell growth and survival critically rely on adequate oxygen supply, hypoxia provides physiologic selective pressure for the expansion of cells with enhanced hypoxia tolerance (5). Many studies have demonstrated that hypoxia selection of death-resistant tumor cells involves the regulation of p53, a tumor suppressor gene, and defects in the cell death

mechanism, such as the upregulation of antiapoptotic proteins (6–11).

Macrophages, cells present in diseased tissues, display functional and phenotypic heterogeneity (12, 13). Moreover, macrophages adapt metabolically and energetically to reduced oxygen tension through the glycolytic processing of glucose (2, 14). Recent studies have shown that in macrophages, hypoxia alters viability, metabolic and phagocytic activities, the release of cytokines, and susceptibility to microorganism infections (15–18). Most importantly, Yun *et al.* (19) demonstrated that subpopulations of macrophages resistant to chronic hypoxia exposure display differential phenotypic characteristics with respect to low-density lipoprotein (LDL) uptake, heat-shock protein 70 kDa (HSP70) expression, and tumor necrosis factor- α (TNF- α) production. The authors suggested that macrophages that are adapted to the hypoxic stress of atherosclerotic lesions may play a critical role in the perpetuation of inflammatory vascular sites (19).

Our laboratory has been interested in the influence of low oxygen tension on macrophage response to parasite stimulus (16, 20, 21). Macrophages treated with a single exposure to mild hypoxia (6% O₂) and infected with *Leishmania amazonensis* were able to reduce intracellular parasitism (16, 20). This protozoan is an intramacrophage parasite that causes chronic human diseases ranging from localized to diffuse cutaneous infections (22, 23). The former is the most common form of leishmaniasis and is characterized by a skin ulcer presenting a raised, expanded border of parasite-infected macrophages with an incomplete granulomatous reaction and a necrotic tissue center (22, 24). These lesions self-heal over months or years (22). Diffuse cutaneous leishmaniasis is associated with cutaneous metastases from the initial skin lesion (22, 23). Since chronic infectious diseases, such as leishmaniasis, result in long-term hypoxic tissue or repeated hypoxia stress (21, 25), the occurrence of resistant macrophages by means of hypoxia selection is highly likely. The selection of such cells may contribute to the control or chronicity of leishmaniasis and interfere with the treatment outcome. In this study, a mouse macrophage cell line (J774 cells) was used to obtain subpopulations of death-resistant macrophages induced by long-term exposure to severe hypoxia

This work was supported by Fundação de Amparo a Pesquisa do Estado de São Paulo and Conselho Nacional de Desenvolvimento Científico e Tecnológico.

¹ To whom correspondence should be addressed at Department of Parasitology, Biology Institute, Universidade Estadual de Campinas, Caixa Postal 6109, Cep 13083-970, Campinas, São Paulo, Brazil. E-mail: sgiorgio@unicamp.br

Received April 5, 2006.
Accepted May 29, 2006.

1535-3702/07/2321-0088\$15.00
Copyright © 2007 by the Society for Experimental Biology and Medicine

(<1% O₂). These hypoxia-selected macrophages were evaluated with regard to HSP70 expression, TNF- α secretion, nitric oxide (NO) production, and parasiticidal activity.

Materials and Methods

Reagents. All reagents were purchased from Sigma Chemical Co. (St. Louis, MO), except recombinant murine TNF- α (BD Biosciences, San Diego, CA); anti-hypoxic-induced transcription factor 1 α (anti-HIF-1 α) antibody (Santa Cruz Biotechnology Inc., Santa Cruz, CA); fetal calf serum (FCS) (Cultilab, Campinas, Brazil); cell culture plates, flasks, and glass coverslips (Nalge Nunc International, Roskilde, Denmark); glycerol; 2-mercaptoethanol; ethylenediaminetetraacetic acid; pepstatin; phenylmethylsulfonyl fluoride; and nitrocellulose membranes (Amersham Pharmacia Biotech, Piscataway, NJ).

Cell Cultures and Parasite. RPMI 1640 medium was reconstituted according to the manufacturer's instructions and was supplemented with 25 μ g/ml gentamicin, 2 mM L-glutamine, 100 mM HEPES, and 10% FCS. The murine macrophage cell line J774 obtained from the American Type Culture Collection (Manassas, VA) and the murine fibroblast cell line L929 obtained from Banco de Células do Rio de Janeiro (Rio de Janeiro, Brazil) were maintained in RPMI 1640 medium as previously described (26, 27). *Leishmania amazonensis* (MHOM/BR/73/M2269) amastigotes were isolated from active skin lesions of BALB/c mice as previously described (28). The parasites were suspended in RPMI 1640 medium and used immediately after isolation.

Hypoxic Selection of J774 Macrophages. J774 macrophages cultured in RPMI 1640 medium were placed into a gas-tight modular chamber (Billups-Rothenberg, Del Mar, CA). The chamber was gassed for at least 15 mins at a flow rate of 2 liters per minute using certified gases containing CO₂ and N₂ (White Martins Gases, Rio de Janeiro, Brazil), and placed in a 37°C temperature-controlled incubator. The percentage of O₂ was verified by measuring the outflow of gas at the end of the initial flushing period and then at 48-hr intervals using a Fyrite apparatus (Bacharach Inc., Pittsburgh, PA). In all experiments, cell exposure to <1% O₂, 5% CO₂, and balanced N₂ is referred to as hypoxia, and cell exposure to 21% O₂, 5% CO₂, and balanced N₂ is referred to as normoxia. The oxygen tension in the culture medium under hypoxia conditions was 7 mm Hg, and it was 150 mm Hg under normoxia conditions (O₂ analyzer YSI/53; Yellow Springs Instruments Inc., Yellow Springs, OH). The pH of the medium was 7.4 and did not change significantly during the course of the experiments. After 48 hrs of hypoxia exposure, detached J774 macrophages were removed with the RPMI 1640 medium. The remaining attached cells were then cultured in fresh medium and exposed again to hypoxia for 48 hrs. This procedure was repeated five times with the

attached cells that survived each round of hypoxia treatment. After five hypoxic cycles, selected macrophages were immediately used for experiments (Jsel) and also were cultured under normoxic conditions for 15 days (Jsel-15). Cell viability was determined by trypan blue exclusion, ability of adherence, and cell growth (16).

Macrophage Activation and Infection with *L. amazonensis*. For the macrophage activation experiments, the cells were treated with 20 ng/ml mouse interferon γ (IFN- γ) and 10 ng/ml lipopolysaccharide (LPS) from *Escherichia coli* for 24 hrs (26). J774 macrophages were infected by adding to the cell cultures a suspension of living *L. amazonensis* amastigotes in RPMI 1640 medium with a 3-fold excess of parasites for 24 hrs. After the infection period, the cultures were washed to remove extracellular parasites. For the evaluation of the infection index (percentage of infected macrophages \times number of amastigotes per macrophage), cells on coverslips were stained with Giemsa and examined microscopically at \times 1000 magnification (28, 29).

TNF- α Bioassay. The culture supernatants were assayed for TNF- α in a cytotoxicity assay using L929 cells pretreated with 5 μ g/ml actinomycin D in 96-well microtiter plates (27, 30). The sensitivity of the L929 cell line was determined by using recombinant murine TNF- α . After 24 hrs of incubation, the plates were fixed in 20% methanol and stained with 0.5% crystal violet before cytotoxicity quantitation with an automated micro-ELISA reader (Lab-system Multiskan MS, Helsinki, Finland). The cytotoxicity effects by supernatants containing TNF- α activity were calculating using the following formula: cytotoxicity (%) = [OD₅₄₀ control - OD₅₄₀ test] / OD₅₄₀ control \times 100, where control and test represent absorption of L929 plus medium and L929 plus supernatants, respectively (30).

Nitrite Assay. A colorimetric assay measured nitrite content in the supernatant of macrophage cultures (31). Supernatants were incubated with Griess reagent, and the absorbance at 540 nm was determined. Nitrite quantities were calculated from a standard curve with serial dilutions of sodium nitrite in RPMI 1640 medium. Each sample was assayed in triplicate, and the results are mean values \pm SD.

Immunoblotting Analyses. After different treatments, the macrophages were scraped from the culture flasks, checked for viability, and then rinsed twice with phosphate-buffered saline. Lysis buffer (62.5 mM Tris-HCl, pH 6.8; 69 mM sodium dodecyl sulfate [SDS]; 10% glycerol; 2% 2-mercaptoethanol; 34 mM ethylenediaminetetraacetic acid; 2 μ g/ml pepstatin; and 1 mM phenylmethylsulfonyl fluoride) was added to the cell pellets. Proteins were denatured at 95°C for 3 mins, electrophoresed on a 10% SDS-polyacrylamide gel system (Thermo EC, Holbrook, NY) and transferred to nitrocellulose membranes. After blotting, the membranes were incubated with mouse monoclonal anti-HSP70, anti-inducible nitric oxide synthase (anti-iNOS), or rabbit polyclonal anti-HIF-1 α antibodies. The secondary antibody consisted of peroxidase-




Figure 1. Western blot analyses of HSP70 in J774 and hypoxia-selected macrophages. (A) J774 cells cultured in normoxia for 24 hrs (lane 1), J774 cells cultured in hypoxia for 24 hrs (lane 2), Jsel cells cultured in normoxia for 24 hrs (lane 3), Jsel cells cultured in hypoxia for 24 hrs (lane 4). Graph shows the results of Western blots analyzed by densitometry. (B) J774 cells cultured in normoxia for 24 hrs (lane 1), J774 cells cultured in hypoxia for 24 hrs (lane 2), Jsel-15 cells cultured in normoxia for 24 hrs (lane 3), Jsel-15 cells cultured in hypoxia for 24 hrs (lane 4). Graph shows the results of Western blots analyzed by densitometry.

conjugated rabbit anti-mouse IgG or goat anti-rabbit IgG; development was performed with 3,3-diaminobenzidine. Immunoreaction images were scanned, and the densitometric value of each band was determined using Image Master Total Lab version 1 software (Amersham Pharmacia Biotech; Ref. 20).

Statistical Evaluation. The results are expressed as mean \pm SD. Data obtained from the experiments with J774, Jsel, and Jsel-15 macrophages were analyzed statistically by the Student *t*-test, with a significance level set at $P < 0.05$ or $P < 0.01$.

Results

Selection of Hypoxia-Induced Death-Resistant Macrophages. Before any experiments of hypoxic selection, the induction of biologically relevant hypoxia in our cell system was confirmed by immunoblotting analyses of HIF-1 α expression in J774 macrophages exposed to severe hypoxia (<1% O₂) for 24 hrs. In contrast, HIF-1 α could not be detected in J774 macrophages maintained in normoxia (data not shown). Our strategy was to subject J774 macrophages to hypoxia to kill vulnerable cells and select resistant macrophages. To this end, macrophages were exposed to hypoxia for 48 hrs, and after this first hypoxic cycle the detached cells were removed and the remaining attached macrophages were again exposed to hypoxia. This procedure was repeated five times with the attached cells that survived each round of hypoxia exposure. The cell viability during the first hypoxic cycle was about 40%. The macrophage viability rate increased to about 80% and 90% after three and five sequential hypoxic cycles, respectively, demonstrating that the selected macrophages became resistant to death induced by hypoxia. No differences in morphology, proliferation, and adhering ability were

observed between J774 macrophages and hypoxia-selected macrophages (Jsel; data not shown).

Expression of HSP70 by Hypoxia-Selected Macrophages. HSP70 is a heat-shock protein expressed constitutively in macrophages, and previous studies indicated a reduction in HSP70 expression level in cells after a single period (24 hrs) of mild hypoxia ($\geq 6\%$ O₂; Refs. 20, 32, 33). To see whether similar changes in HSP70 expression were detectable in hypoxia-selected macrophages, Western blots of HSP70 were performed. As shown in Figure 1, HSP70 was constitutively expressed in J774 macrophages in normoxia, and a 40% reduction in HSP70 was observed after exposure to hypoxia (<1% O₂) (Fig. 1A, lanes 1 and 2). Jsel macrophages maintained in normoxia for 24 hrs expressed HSP70 at lower levels than J774 cells cultivated in normoxia (Fig. 1A, lanes 1 and 3). In contrast to J774 macrophages, the level of HSP70 in Jsel cells was not reduced after 24 hrs of hypoxia (Fig. 1, lanes 3 and 4). Notably, the HSP70 expression pattern of Jsel cells was maintained in this macrophage subpopulation, even after 15 days of cell culture under normoxia conditions (Jsel-15); that is, they showed reduced HSP70 expression compared with J774 macrophages cultivated in normoxia (Fig. 1B, lanes 1 and 3) and increased HSP70 expression when cultured for 24 hrs in hypoxia (Fig. 1B, lanes 3 and 4). These data indicate that hypoxia selection of a distinct HSP70-related phenotype was not a transient phenomenon but was retained in macrophages for at least 2 weeks under ambient O₂ conditions.

Production of TNF- α and NO by Hypoxia-Selected Macrophages. We further examined whether the selection of hypoxia-resistant macrophages affected the ability to produce two biomolecules, TNF- α and NO, involved in inflammatory responses (34, 35). J774, Jsel, and




Figure 2. TNF- α production of J774 and hypoxia-selected macrophages. TNF- α bioactivity from culture supernatants of J774, Jsel, and Jsel-15 cells exposed to normoxia (white bars) or hypoxia (black bars), or stimulated with IFN- γ + LPS in normoxia (gray bars) for 24 hrs, were determined in the L929 cytotoxicity assay. The results presented were obtained in triplicate L929 cultures \pm SD of one representative experiment. * $P < 0.01$ compared with culture supernatants of J774 cells in normoxia.

Jsel-15 macrophages were cultured under normoxia or hypoxia conditions or were stimulated with IFN- γ plus LPS, a well-defined proinflammatory activator system (35, 36), and 24 hrs later supernatants were collected and assayed for TNF- α bioactivity. As shown in Figure 2, a significant difference in TNF- α release between J774, Jsel, and Jsel-15 release was observed. The highest TNF- α bioactivity was observed in supernatant cultures of IFN- γ + LPS-exposed J774 cells (Fig. 2). The Jsel macrophage cultures presented a significantly lower TNF- α release than J774 and Jsel-15 cell cultures when treated with the same stimulus (IFN- γ +

LPS). When the three macrophage populations were exposed to 24 hrs of hypoxia, TNF- α bioactivity was enhanced in J774 and Jsel supernatants (Fig. 2). Notably, hypoxia reduced TNF- α bioactivity in Jsel-15 supernatant cultures (Fig. 2).

To examine whether the selection of hypoxia-resistant macrophages also affected their ability to produce NO, the macrophages were treated with IFN- γ + LPS and assayed for nitrite accumulation in the supernatants (Fig. 3A). As previously reported, high levels of nitrite were induced in J774 macrophage cultures stimulated with IFN- γ + LPS in normoxic conditions (26). In contrast to J774 cells, Jsel and Jsel-15 macrophages produced low levels of nitrite in response to the activators (about 30% of J774 macrophage nitrite accumulation; Fig. 3A). It should be noted that the three subpopulations of macrophages (J774, Jsel, and Jsel-15 cells) produced minimum quantities of nitrite under hypoxic condition (data not shown). The question of whether increased nitrite production during activation with IFN- γ + LPS in normoxia is related to increased iNOS expression in macrophages was addressed by measuring the levels of iNOS protein in cell lysates. As shown in Figure 4B, J774 cells treated with IFN- γ + LPS showed a major increase in iNOS protein levels (Fig. 3B, lane 1). iNOS also increased after IFN- γ + LPS stimulation in Jsel and Jsel-15 macrophages; however, the iNOS protein level was about 40% lower than in J774 cells (Fig. 3B and C). There was no detectable iNOS protein in nonstimulated macrophages cultured under normoxia or hypoxia (data not shown). Thus, a differential response of hypoxia-selected macrophages to inflammatory stimulus was demonstrated by the differences




Figure 3. NO production of J774 and hypoxia-selected macrophages. (A) J774, Jsel, and Jsel-15 macrophages were stimulated with IFN- γ + LPS in normoxia for 24 hrs. Nitrite concentrations were determined in the supernatants from cultures by the Griess reaction. The results represent the mean \pm SD of one representative experiment. * $P < 0.05$ compared with culture supernatants of J774 cells stimulated with IFN- γ + LPS. (B) Western blot analyses of iNOS in J774 (lane 1) and Jsel (lane 2). Graph shows the results of Western blots analyzed by densitometry. (C) Western blot analyses of iNOS in J774 (lane 1) and Jsel-15 (lane 2). Graph shows the results of Western blots analyzed by densitometry.




Figure 4. Leishmania infection in J774 and hypoxia-selected macrophages. J774, Jsel, and Jsel-15 macrophages were infected with *Leishmania* amastigotes and cultured in normoxia (white bars) or hypoxia (black bars). After 24 hrs the IF was determined as described in Materials and Methods. The results represent the mean \pm SD of one representative experiment. * $P < 0.05$; ** $P < 0.01$ compared with J774 macrophages infected with *Leishmania* amastigotes and cultured in normoxia.

in the levels of TNF and NO production observed between J774, Jsel, and Jsel-15 cells.

Susceptibility of Hypoxia-Selected Macrophages for an Intracellular Pathogen. To address the question of whether the selection of hypoxia-resistant macrophages affected functional ability, the susceptibility of the cells to an intracellular pathogen, *Leishmania*, was analyzed. Macrophages are the host cells for *Leishmania*, where these parasites are able to survive and replicate (37). As expected, J774 macrophages cultured with *Leishmania* and maintained under normoxia for 24 hrs showed a high infection index (IF) (Fig. 4). However, in J774 cell cultures subjected to hypoxia, a reduction in IF occurred (Fig. 4). Notably, Jsel macrophages exposed to normoxia during the period of infection showed a significant reduction in infection compared with J774 cells (Jsel IF: 65; J774 IF: 250; Fig. 4). A more pronounced reduction in IF was observed when Jsel macrophages were infected under hypoxia (IF: 24, Fig. 4). The ability of hypoxia-selected macrophages to resist *Leishmania* infection also was maintained in Jsel-15 cells. IF was significantly suppressed in Jsel-15 cells exposed to normoxia and hypoxia during the period of infection (Fig. 4). These results indicate that hypoxia-selected macrophages are more resistant to an intracellular pathogen infection.

Discussion

It is clear from this and previous studies (6–11, 19) that hypoxic-rendered cell populations show death resistance to microenvironments with low oxygen pressure. Altered characteristics in tumor cells subjected to selective pressure, such as hypoxia, have been reported, and the selected cells

generally are examined by growth characteristic, resistance to apoptotic stimulus, and response to DNA-damaging drugs (9–11). The aim of the present study was to investigate whether death-resistant macrophages induced by hypoxia selection were accompanied by functional changes. The J774 macrophage cell line was used to obtain two subpopulations of death-resistant macrophages induced by long-term exposure to severe hypoxia ($<1\%$ O_2): Jsel, assayed immediately after five cycles of hypoxia; and Jsel-15, derived from Jsel macrophages and cultured for 15 days in normoxia before use. We found that HSP70, a protein that has been correlated with cell resistance to micro-environment stress (38), was differentially expressed in J774, Jsel, and Jsel-15 macrophages. The expression level of HSP70 was reduced in J774 macrophages exposed to hypoxia, whereas the HSP70 level was not altered in Jsel macrophages. Jsel-15 cells express low levels of HSP70 in normoxia and high levels of HSP70 in hypoxia. Yun *et al.* (19) also noted that the RAW macrophage cell line exposed to hypoxia (2% O_2) expressed low levels of HSP70, and selected macrophages maintained the same HSP70 levels under hypoxia or normoxia. The present results agree with their findings and indicate that low levels of HSP70 in macrophages coincide with poor resistance to severe hypoxia. Since the present work examined a subpopulation of macrophages cultured for 15 days in normoxia before the tests (Jsel-15), which showed enhanced levels of HSP70 after hypoxic stimulus, it is possible to confirm that HSP70 is a marker for adaptation to severe hypoxic stress.

The present study also demonstrated macrophage variability in the pattern of TNF- α and NO production. These two biomolecules are involved in macrophage cytotoxic and inflammatory responses (34, 35). The results demonstrated maximum TNF- α bioactivity in J774 macrophage cultures after IFN- γ + LPS stimulation, an intermediate TNF- α production by cells cultured under hypoxia, and low TNF- α production by nonstimulated cells. In Jsel macrophages, an intermediate TNF- α production was observed in cultures stimulated with IFN- γ + LPS or maintained in hypoxia. In contrast, Jsel-15 cells produced maximum TNF- α levels when stimulated with IFN- γ + LPS, but they did not respond to a hypoxic microenvironment. These results support previous observations that hypoxia (1%–9% O_2) induces TNF- α release in unstimulated or endotoxin-stimulated human mononuclear cells and macrophage cell lines (27, 39–42). However, Yun *et al.* (19) reported a suppressive effect of hypoxia (2% O_2) on the murine macrophage cell line RAW 264.7, although hypoxia-selected RAW macrophages secreted TNF- α when exposed to hypoxia (19). Thus, despite the different results achieved in experiments with these two parental macrophages (J774 and RAW cells), which likely reflect differences in cell line type, the present results with hypoxia-selected macrophages are in agreement with those by Yun *et al.* (19), indicating a distinct phenotype for TNF- α production in hypoxia-selected macrophages. The current

experiments extend the studies by Yun *et al.* (19) and reinforce the hypothesis that a distinct functional phenotype is originated after hypoxia selection. Hypoxia-selected macrophages also differ from J774 macrophages with respect to NO production. The present results indicated that under normoxic conditions, IFN- γ + LPS-activated Jsel and Jsel-15 macrophages produce less nitrite and iNOS protein than IFN- γ + LPS-activated J774 macrophages. Although no reports on the effect of hypoxia selection on NO production by selected cells are available for comparison with these results, a number of studies have reported that nitrite accumulation and iNOS protein in macrophages and myocytes are decreased in proportion to oxygen tension (42–46). The absence of oxygen as a substrate (43, 45), an increase in cysteine protease calpain expression contributing to iNOS degradation (47), and a lack of tetrahydrobiopterin cofactor and auxiliary proteins (43, 48), resulting in the displacement and instability of the iNOS structure (49), are used to explain low NO production in cells exposed to hypoxia. Whether transcriptional or posttranscriptional events are involved in decreased NO and iNOS production in hypoxia-selected macrophages remains to be established.

To evaluate whether another functional marker was altered in hypoxia-selected macrophages, the ability to maintain or endure infection by an intracellular parasite, *Leishmania*, was analyzed in J774, Jsel, and Jsel-15 cells. Macrophages are hosts for *Leishmania* and are the major site for parasite replication (37). The results demonstrated that hypoxia caused a significant decrease in the J774 macrophage IF by *Leishmania*. More interestingly, we found that hypoxia-selected macrophages are resistant to parasites even under normoxic conditions, indicating that hypoxia selection confers a *Leishmania*-resistant phenotype to macrophages. To our knowledge no previous studies have investigated the effect of cell hypoxia selection with regard to microorganism infection. Yun *et al.* (19) demonstrated that hypoxia-selected RAW macrophages take up a minimum level of acetylated LDL in normoxia (19). Although direct comparison with these results is not possible, collectively they indicate deficient phagocytosis in hypoxia-selected macrophages, which could be the effect of alterations in cytoskeleton protein rearrangement, changes in ligands and receptors, and membrane trafficking (19, 49, 50). In support of this interpretation, a number of studies have reported the distinct changes in gene and protein expression that occur in macrophages and other cell types when they experience *in vitro* hypoxia (49, 51–53). These include the modulation of molecules required for metabolism, transcriptional factors, and membrane-bound proteins (e.g., CD80) and proteins associated with signaling cascade (17, 41, 49, 51–53).

In summary, this study shows that exposing macrophages to periods of severe hypoxia results in the selection of cells with phenotypes associated with the modulation of HSP70 expression, TNF- α and NO production, and *Leishmania* susceptibility. This variability in the pattern of

HSP70 expression, TNF- α and NO production, and parasite susceptibility in hypoxia-selected and nonselected J774 macrophage subpopulations suggest that macrophages can progressively change their phenotype in response to progressive changes in microenvironmental signals. Many questions remain concerning the complex molecular mechanism that alters macrophage phenotype after hypoxic selection. HIF-1 is a heterodimer consisting of an alpha subunit, the oxygen-responsive component, and beta subunit constitutively expressed in most cells (55). HIF-1 α induced under hypoxic conditions regulates the expression of several genes, including glycolytic enzymes and cytokines (55). It has recently been shown that during functional maturation from monocytes to macrophages HIF-1 activity increases, and cells acquire the ability to respond to hypoxia more robustly (56). Since HIF-1 α expression could be detected in J774 macrophages under hypoxic conditions as chosen for our experiments, we can speculate that HIF-1 α is involved in the phenotype and functional changes observed in hypoxia-selected macrophages.

Considering the presence of hypoxia in inflammatory and infectious lesions from initiation to resolution, it is likely that injured tissues would contain hypoxia-selected macrophages with the attenuated inflammatory phenotype, which would facilitate tissue regeneration and homeostasis. Therefore, knowledge regarding the functional state and plasticity of macrophages (54) challenged by the tissue microenvironment could be of paramount importance for drug and vaccine design.

1. Blouw B, Song H, Tihan T, Bosze J, Ferrara N, Gerber HP, Johnson RS, Bergers G. The hypoxic response of tumors is dependent on their microenvironment. *Cancer Cell* 4:133–146, 2003.
2. Albina JE, Reichner JS. Oxygen and the regulation of gene expression in wounds. *Wound Repair Regen* 11:445–451, 2003.
3. Simanionok JP. Non-ischemic hypoxia of the arterial wall is a primary cause of atherosclerosis. *Med Hypotheses* 46:155–161, 1996.
4. Ikeda E. Cellular response to tissue hypoxia and its involvement in disease progression. *Pathol Int* 55:603–610, 2005.
5. Yun JK, McCormick TS, Judware R, Lapetina EG. Cellular adaptive responses to low oxygen tension: apoptosis and resistance. *Neurochem Res* 22:517–521, 1997.
6. Honsik CJ, Diamant M, Olsson L. Generation of stable cellular phenotypes in a human malignant cell line conditioned by alterations in the cellular microenvironment. *Cancer Res* 46:940–949, 1986.
7. Shimizu S, Eguchi Y, Kamiike W, Itoh Y, Hasegawa J, Yamabe K, Otsuki Y, Matsuda H, Tsujimoto Y. Induction of apoptosis as well as necrosis by hypoxia and predominant prevention of apoptosis by Bcl-2 and Bcl-XL. *Cancer Res* 56:2161–2166, 1996.
8. Graeber TG, Osmanian C, Jacks T, Housman DE, Koch CJ, Lowe SW, Giaccia AJ. Hypoxia-mediated selection of cells with diminished apoptotic potential in solid tumours. *Nature* 379:88–91, 1996.
9. Weinmann M, Jendrossek V, Guner D, Goecke B, Belka C. Cyclic exposure to hypoxia and reoxygenation selects for tumor cells with defects in mitochondrial apoptotic pathways. *FASEB J* 18:1906–1908, 2004.
10. Dong Z, Wang J. Hypoxia selection of death-resistant cells. A role for Bcl-X(L). *J Biol Chem* 279:9215–9221, 2004.

11. Yao K, Gietema JA, Shida S, Selvakumaran M, Fonrose X, Haas NB, Testa J, O'Dwyer PJ. In vitro hypoxia-conditioned colon cancer cell lines derived from HCT116 and HT29 exhibit altered apoptosis susceptibility and a more angiogenic profile in vivo. *Br J Cancer* 93: 1356–1363, 2005.
12. DiPietro LA, Strieter RM. Macrophages in wound healing. In: Burke B, Lewis CE, Eds. *The Macrophage*. Oxford, UK: Oxford University Press, p435, 2002.
13. Hamilton TA. Molecular basis of macrophage activation. In: Burke B, Lewis CE, Eds. *The Macrophage*. Oxford, UK: Oxford University Press, p73, 2002.
14. Simon LM, Axline SG, Horn BR, Robin ED. Adaptations of energy metabolism in the cultivated macrophage. *J Exp Med* 138:1413–1425, 1973.
15. Lewis JS, Lee JÁ, Underwood JC, Harris AL, Lewis CE. Macrophage responses to hypoxia: relevance to disease mechanisms. *J Leukoc Biol* 66:889–900, 1999.
16. Colhone MC, Arrais-Silva WW, Picoli C, Giorgio S. Effect of hypoxia on macrophage infection by *Leishmania amazonensis*. *J Parasitol* 90: 510–515, 2004.
17. Murdoch C, Muthana M, Lewis CE. Hypoxia regulates macrophage functions in inflammation. *J Immunol* 175:6257–6263, 2005.
18. Peyssonnaud C, Datta V, Cramer T, Doedens A, Theodorakis EA, Gallo RL, Hurtado-Ziola N, Nizet V, Johnson RS. HIF-1 α expression regulates the bactericidal capacity of phagocytes. *J Clin Invest* 115:1806–1815, 2005.
19. Yun JK, McCormick TS, Villabona C, Judware RR, Espinosa MB, Lapetina EG. Inflammatory mediators are perpetuated in macrophages resistant to apoptosis induced by hypoxia. *Proc Natl Acad Sci U S A* 94:13903–13908, 1997.
20. Degrossoli A, Colhone MC, Arrais-Silva WW, Giorgio S. Hypoxia modulates expression of the 70-kD heat shock protein and reduces *Leishmania* infection in macrophages. *J Biomed Sci* 11:847–854, 2004.
21. Arrais-Silva WW, Paffaro VA Jr, Yamada AT, Giorgio S. Expression of hypoxia-inducible factor-1 α in the cutaneous lesions of BALB/c mice infected with *Leishmania amazonensis*. *Exp Mol Pathol* 78:49–54, 2005.
22. Grimaldi G Jr, Tesh RB. Leishmaniasis of the New World: current concepts and implications for future research. *Clin Microbiol Rev* 6: 230–250, 1993.
23. Desjeux P. Leishmaniasis: current situation and new perspectives. *Comp Immunol Microbiol Infect Dis* 27:305–318, 2004.
24. Moll H, Ritter U, Flohe S, Erb K, Bauer C, Blank C. Cutaneous leishmaniasis: a model for analysis of the immunoregulation by accessory cells. *Med Microbiol Immunol (Berl)* 184:163–168, 1996.
25. Kanan MW. Mucocutaneous leishmaniasis in guinea-pigs inoculated intravenously with *Leishmania enriettii*. *Br J Dermatol* 92:663–673, 1975.
26. Linares E, Augusto O, Barão SC, Giorgio S. *Leishmania amazonensis* infection does not inhibit systemic nitric oxide levels elicited by lipopolysaccharide in vivo. *J Parasitol* 86:78–82, 2000.
27. Leeper-Woodford SK, Detmer K. Acute hypoxia increases alveolar macrophage tumor necrosis factor activity and alters NF- κ B expression. *Am J Physiol* 276:L909–L916, 1999.
28. Barbieri CL, Giorgio S, Merjan AJ, Figueiredo EN. Glycosphingolipid antigens of *Leishmania (Leishmania) amazonensis* amastigotes identified by use of a monoclonal antibody. *Infect Immun* 61:2131–2137, 1993.
29. Pereira WK, Lonardoni MV, Grespan R, Caparroz-Assef SM, Cuman RK, Bersani-Amado CA. Immunomodulatory effect of Canova medication on experimental *Leishmania amazonensis* infection. *J Infect* 51:157–164, 2004.
30. Moore K, Thomas A, Harding KG. Iodine released from the wound dressing Iodosorb modulates the secretion of cytokines by human macrophages responding to bacterial lipopolysaccharide. *Int J Biochem Cell Biol* 29:163–167, 1997.
31. Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JS, Tannenbaum SR. Analysis of nitrate, nitrite, and [¹⁵N] nitrate in biological fluids. *Anal Biochem* 126:131–138, 1982.
32. Oehler R, Schmierer B, Zellner M, Prohaska R, Roth E. Endothelial cells downregulate expression of 70 kDa heat shock protein during hypoxia. *Biochem Biophys Res Commun* 274:542–547, 2000.
33. Mohan RM, Golding S, Paterson DJ. Intermittent hypoxia improves atrial tolerance to subsequent anoxia and reduces stress protein expression. *Acta Physiol Scand* 172:89–95, 2001.
34. Kollias G. TNF pathophysiology in murine models of chronic inflammation and autoimmunity. *Semin Arthritis Rheum* 34:3–6, 2005.
35. Korhonen R, Lahti A, Kankaanranta H, Moilanen E. Nitric oxide production and signaling in inflammation. *Curr Drug Targets Inflamm Allergy* 4:471–479, 2005.
36. Stuehr DJ, Marletta MA. Induction of nitrite/nitrate synthesis in murine macrophages by BCG infection, lymphokines, or interferon- γ . *J Immunol* 139:518–525, 1987.
37. Handman E, Bullen DV. Interaction of *Leishmania* with the host macrophage. *Trends Parasitol* 18:332–334, 2002.
38. Kiang JG, Tsokos GC. Heat shock protein 70 kDa: molecular biology, biochemistry, and physiology. *Pharmacol Ther* 80:183–201, 1998.
39. Scannell G, Waxman K, Kaml GJ, Ioli G, Gatanaga T, Yamamoto R, Granger GA. Hypoxia induces a human macrophage cell line to release tumor necrosis factor- α and its soluble receptors in vitro. *J Surg Res* 54:281–285, 1993.
40. Ghezzi P, Dinarello CA, Bianchi M, Rosandich ME, Repine JE, White CW. Hypoxia increases production of interleukin-1 and tumor necrosis factor by human mononuclear cells. *Cytokine* 3:189–194, 1991.
41. Lahat N, Rahat MA, Ballan M, Weiss-Cerem L, Engelmayer M, Bitterman H. Hypoxia reduces CD80 expression on monocytes but enhances their LPS-stimulated TNF- α secretion. *J Leukoc Biol* 74:197–205, 2003.
42. Ndengele MM, Bellone CJ, Lechner AJ, Matuschak GM. Brief hypoxia differentially regulates LPS-induced IL-1 β and TNF- α gene transcription in RAW 264.7 cells. *Am J Physiol Lung Cell Mol Physiol* 278:L1289–L1296, 2000.
43. Otto CM, Baumgardner JE. Effect of culture PO₂ on macrophage (RAW 264.7) nitric oxide production. *Am J Physiol Cell Physiol* 280: C280–C287, 2001.
44. Kacimi R, Long CS, Karlner JS. Chronic hypoxia modulates the interleukin-1 β -stimulated inducible nitric oxide synthase pathway in cardiac myocytes. *Circulation* 96:1937–1943, 1997.
45. McCormick CC, Li WP, Calero M. Oxygen tension limits nitric oxide synthesis by activated macrophages. *Biochem J* 350:709–716, 2000.
46. Frost MT, Wang Q, Moncada Singer M. Hypoxia accelerates nitric oxide-dependent inhibition of mitochondrial complex I in activated macrophages. *Am J Physiol Regul Integr Comp Physiol* 288:R394–R400, 2005.
47. Su Y, Block ER. Role of calpain in hypoxic inhibition of nitric oxide synthase activity in pulmonary endothelial cells. *Am J Physiol Lung Cell Mol Physiol* 278:L1204–L1212, 2000.
48. Stuehr DJ, Wei CC, Wang Z, Hille R. Exploring the redox reactions between heme and tetrahydrobiopterin in the nitric oxide synthases. *Dalton Trans* 21:3427–3435, 2005.
49. Daniliuc S, Bitterman H, Rahat MA, Kinarty A, Rosenzweig D, Lahat N. Hypoxia inactivates inducible nitric oxide synthase in mouse macrophages by disrupting its interaction with α -actinin 4. *J Immunol* 171:3225–3232, 2003.
50. Aderem A, Underhill DM. Mechanisms of phagocytosis in macrophages. *Annu Rev Immunol* 17:593–623, 1999.
51. Weinmann M, Belka C, Guner D, Goecke B, Muller I, Bamberg M, Jendrosseck V. Array-based comparative gene expression analysis of

- tumor cells with increased apoptosis resistance after hypoxic selection. *Oncogene* 24:5914–5922, 2005.
52. Kumar GK, Klein JB. Analysis of expression and posttranslational modification of proteins during hypoxia. *J Appl Physiol* 96:1178–1186, 2004.
53. Ning W, Chu TJ, Li CJ, Choi AM, Peters DG. Genome-wide analysis of the endothelial transcriptome under short-term chronic hypoxia. *Physiol Genomics* 18:70–78, 2004.
54. Stout RD, Suttles J. Functional plasticity of macrophages: reversible adaptation to changing microenvironments. *J Leukoc Biol* 76:509–513, 2004.
55. Semenza GL. Hypoxia-inducible factor 1: oxygen homeostasis and disease pathophysiology. *Trends Mol Med* 7:345–350, 2001.
56. Oda T, Hirota K, Nishi K, Takabuchi S, Oda S, Yamada H, Arai T, Fukuda K, Kita T, Adachi T, Semenza GL, Nohara R. Activation of hypoxia-inducible factor 1 during macrophage differentiation. *Am J Physiol Cell Physiol* 291:C104–C113, 2006.